

Recommended Childhood Immunization Schedule Utah Valley Pediatrics

| Vaccine | Birth | 2 months | 4 months | 6 months | 12 months | 15-18 months | 2 years | 5 years | 11-15 years |
|--------------------------------|-------|-----------|-----------|---------------------------------|-----------|--------------------|---------------|-----------|------------------|
| Hepatitis B | Hep B | Hep B | | Hep B | | | | | |
| Diphtheria, Tetanus, Pertussis | | DTaP | DTaP | DTaP | | DTaP | | DTaP | Tdap |
| Haemophilus influenzae type b | | Hib | Hib | Hib | | Hib | | | |
| Inactivated Poliovirus | | IPV | IPV | IPV | | | | IPV | |
| Pneumococcal | | PNC | PNC | PNC | PNC | | | | |
| Mumps, Measles, Rubella | | | | | MMR | | | MMR | |
| Varicella | | | | | Varicella | | | Varicella | |
| Hepatitis A | | | | | Hep A | Hep A ¹ | | | |
| Meningococcal | | | | | | | Meningococcal | | |
| Rotavirus (optional) | | Rotavirus | Rotavirus | Rotavirus | | | | | |
| Human Papillomavirus | | | | | | | | | HPV ² |
| Influenza | | | | Influenza (Yearly) ³ | | | | | |

Recommended number of doses by 2 years: 3 Hep B, 4 DtaP, 4 Hib, 3 IPV, 4 PNC, 1 MMR, 1 Varicella, and 2 Hep A.

Hep A¹: Second dose administered at 18 months or 2 years.

HPV²: Available to females age 9-26 years. Three doses administered 0, 2, and 6 months apart.

Influenza (Yearly)³: Can be administered at 6 months.