



Ibuprofen
(Advil, Motrin, Nuprin, Medipren)

Give every 6 hours for fever over 102° rectal	May Give Every	12-17 lbs. 6-11 mo.	18-23 lbs. 12-23 mo.	24-35 lbs. 2-3 yr.	36-47 lbs. 4-5 yr.	48-59 lbs. 6-8 yr.	60-85 lbs. 9-11 yr.
Chewable Tablets 100 mg	6 hours			1 tab.	1 1/2 tab.	2 tab.	2 1/2 tab.
Chewable Tablets 50 mg	6 hours			2 tab.	3 tab.	4 tab.	5 tab.
DROPS 50 mg/1.25 ml *	6 hours	1 dpr. (1.25 ml.)	1 1/2 dpr. (1.875 ml.)	2 dpr. (2.5 ml.)	3 dpr. (3.75 ml.)		
Suspension 100 mg/5 ml	6 hours	1/2 tsp.	3/4 tsp.	1 tsp.	1 1/2 tsp.	2 tsp.	2 1/2 tsp.
Tablets 200 mg	6 hours					1 tab.	1 1/2 tab.

*** CAUTION:** DO NOT GIVE TO AN INFANT UNDER 6 MONTHS OF AGE. Consult with a physician or pharmacist before combining any medication. Some drugs may contain additional pain relievers. Aspirin should not be given to your child.

