

Vaccine	Birth	2 mo.	4 mo.	6 mo.	12 mo.	15-18 mo.	2 yr.	4-6 yr.	11-15 yr.
Hepatitis B	Hep B	Нер В	-	Нер В	-	-	-	-	-
Diphtheria, Tetanus, Pertussis	-	DTaP	DTaP	DTaP	-	DTaP	-	DTaP	Tdap
Haemophilus influenzae type b	-	Hib	Hib	Hib	-	Hib	-	-	-
Inactivated Poliovirus	-	IPV	IPV	IPV	-	IPV	-	IPV	-
Pneumococcal	-	PCV	PCV	PCV	PC	V	-	-	-
Mumps, Measles, Rubella	-	-	-	-	MMR	-	-	MMR	-
Varicella	-	-	-	-	Varicella	-	-	Varicella	-
Hepatitis A	-	-	-	-	Нер А	Нер	o A ¹	-	-
Meningococcal	-	-	-	-	-	-	-	-	Meningococcal
Rotavirus	-	Rotavirus	Rotavirus	Rotavirus	-	-	-	-	-
Human Papillomavirus	-	-	-	-	-	-	-	-	HPV ²
Influenza	-	-	-			Influe	nza (Yearl	y)3	

Recommended number of doses by 2 years: 3 Hep B, 4 DtaP, 4 Hib, 3 IPV, 4 PCV, 1 MMR, 1 Varicella, and 2 Hep A

Hep A¹: Second dose administered at 18 to 2 years

HPV²: Available ages 9-26 years. Two doses under 15 years of age, 3 doses for those 15 years and up

Influenza (Yearly)3: Second dose administered at 18 to 2 years

www.uvpediatrics.com