

Influenza

Guidelines for Parents



The flu — every child seems to get it at some time or another. But what is the flu? Can it be prevented? Should my child get a flu shot? These are just a few of the most common questions parents have about *influenza* (the flu). The following information will help you understand what influenza is, how to prevent it, and treatments that are now available.

What is the flu?

The flu is an illness caused by a virus. There are three different flu viruses, types A, B, and C. Types A (the most common) and B (usually milder) cause the usual epidemics of the flu. Flu viruses usually strike between December and early April. Because each of the types of flu virus has different strains, every year the flu is slightly different. It can infect people several times during their lifetime.

The flu can last a week or even longer. Your child usually will feel the worst during the first 2 or 3 days and may have any of the following symptoms:

- A sudden fever (usually above 101°F)
- Chills and shakes with the fever
- Extreme tiredness
- Headache and body aches
- Dry, hacking cough
- Sore throat
- Vomiting and belly pain
- Stuffy, runny nose

There are usually no serious complications from the flu. However, sometimes an ear infection, a sinus infection, or pneumonia may develop. Talk to your pediatrician if your child says that his ear hurts, he feels congested in his face and head, his cough persists, or his fever lasts beyond 3 to 4 days.

How the flu is spread

The flu is spread from person to person in the following ways:

- Direct hand-to-hand contact
- Indirect contact (for example, if your child touches an infected surface like a toy or a doorknob and then puts her hand to her own eyes, nose, or mouth)
- Virus droplets being passed through the air from coughing or sneezing

The flu spreads very easily, especially in preschool and school-age children. Adults who spend time with children are exposed easily and can get the disease. The virus usually is transmitted just as symptoms begin or in the first several days of the illness.

Treatment

In children older than 1 year of age, type A influenza can be treated with antiviral agents if given in the first day or two of the illness. This can speed recovery. Under some circumstances, antiviral agents can be taken before exposure to the flu and can prevent illness. This is particularly important for children with serious problems who have not had the flu shot. Antibiotics can

Do not give aspirin to your child for the flu. An increased risk of developing Reye syndrome (an illness that can seriously affect the liver and the brain) is associated with aspirin use during bouts of the flu and many other diseases caused by viruses.

be used to fight bacterial infections but have *no* effect on viruses, including the influenza viruses. Extra bed rest, extra fluids, and light, easy-to-digest meals can also help your child feel better.

If your child is uncomfortable because of fever, acetaminophen in proper doses for age and weight will help him feel better.

Do not give your child over-the-counter cough or cold medicines without checking with your pediatrician.

Prevention

Good hygiene is the best way to prevent the flu from spreading to other family members. If your child has the flu, the following will help prevent its spread:

- Teach your child to cover her mouth and nose with a tissue or her sleeve, but not with her hands, when coughing or sneezing. If your child is old enough, teach her how to blow her nose properly.
- Use facial tissues for runny noses and to catch sneezes. Throw them in the trash after each use.
- Avoid kissing your child on or around the mouth or face, though she will need plenty of hugs while she is sick.
- Make sure everyone washes her hands before and after coming in close contact with someone with the flu.
- Wash dishes and utensils in hot, soapy water or in the dishwasher.
- Do not let children share pacifiers, cups, utensils, washcloths, or towels. Never share toothbrushes.
- Use disposable paper cups in the bathroom and kitchen.
- Disinfect. Viruses can live for more than 30 minutes on doorknobs, toilet handles, countertops, even on toys. Use a disinfectant or soap and hot water to keep these areas clean.
- Do not smoke around your child. Children who are exposed to tobacco smoke cough and wheeze more and have a harder time getting over the flu.

Influenza vaccine

There are safe and effective vaccines to protect against the flu. However, they are mainly recommended for children with health problems that make it risky for them to get the flu. This includes children with the following:

- Heart disease
- Lung disease, including asthma
- Immune problems, such as human immunodeficiency virus (HIV) infection
- Blood diseases
- Cancer

- Chronic kidney disease
- Metabolic diseases, such as diabetes
- Long-term aspirin therapy, such as with rheumatoid arthritis

Children 6 through 23 months are encouraged to get a flu shot each fall, as should everyone in the household. Your pediatrician can recommend what is best for your child.

For children younger than 9 years of age, the vaccine requires two injections, given 1 month apart the first year it is given. After that, only one dose is needed. The best time to get the flu vaccine is in late October to early December before the flu season starts. Vaccination should begin earlier for those needing two shots.

Because the strains of flu are different every year, a new flu vaccine is developed each year as well. The vaccine is made from killed flu viruses and helps the immune system fight the flu. Most children are immune within 2 weeks of getting the vaccine. Side effects are almost always minor and include soreness at the site of the injection and a low-grade fever. The flu shot cannot cause influenza.

Important note: Even though there are few side effects to the vaccine, production of the vaccine involves the use of eggs. If your child has had a serious allergic reaction to eggs or egg products, he should be skin tested before getting the vaccine. If skin testing confirms hypersensitivity, the vaccine usually should not be given.

Scientists are working on the development of a nasal spray flu vaccine. This will be a painless and effective way to protect children from the flu.

Influenza or cold?

Both the flu and colds are caused by viruses and share many symptoms, but there are differences. A child with a common cold usually has a lower fever, a runny nose, and only a small amount of coughing. Children with the flu usually feel much sicker, achy, and miserable. Also, the flu tends to strike more quickly than a cold. Stomach upsets and vomiting are more common with the flu than with a cold. Children who have colds usually have enough energy to play and keep up with their usual day-to-day routines. The flu, on the other hand, keeps most children in bed for several days.

When to call the pediatrician

An older child with the flu usually does not need to see the pediatrician unless the condition becomes more serious. If your child is 3 months of age or younger, however, call your pediatrician if she has a fever. For a child older than 3 months of age who has been exposed to the flu, call your pediatrician if your child experiences any of the following:

- Difficulty breathing
- Blue lips or nails
- A cough that just will not go away (for more than 1 week)
- Pain in the ear
- Continued or new onset of fever after 3 to 4 days of illness

If your child seems extremely sick or his condition does not improve, call your pediatrician.

The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

From your doctor

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

The American Academy of Pediatrics is an organization of 57,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of infants, children, adolescents, and young adults.

American Academy of Pediatrics
PO Box 747
Elk Grove Village, IL 60009-0747
Web site — <http://www.aap.org>

Copyright ©1998, Updated 8/02
American Academy of Pediatrics