

Lead Poisoning: Prevention and Screening



Of all the health problems caused by the environment, lead poisoning is the most preventable. Despite this, almost 1 million children in the United States have elevated levels of lead in their blood. Any child can be at risk for lead poisoning. This brochure has been developed by the American Academy of Pediatrics to inform parents about the risks of lead poisoning and how to prevent it. The brochure also discusses lead screening and treatment for lead poisoning.

How can lead hurt my child?

You may have heard that children can be harmed by the lead in pencils. This is not true. There is no actual lead in pencils and there is no lead in the paint on the outside of pencils.

Children *can* be harmed by lead by:

- Getting lead dust from old paint on their hands or toys and then putting their hands in their mouths
- Breathing in lead dust from old paint
- Eating chips of old paint or dirt that contain lead
- Drinking water from pipes lined or soldered with lead

Once lead enters the body, it travels through the bloodstream and is stored mainly in the bones where it can remain for a lifetime. Very high levels of lead in the body may cause many long-term problems, including:

- Kidney problems
- Anemia
- Hearing loss
- Developmental delays
- Growth problems
- Seizures and coma

Most children with high lead levels in their blood show no obvious symptoms until they reach school age. At that point, some may show learning and behavioral problems.

Where can lead be found?

Lead is most often found in the following places:

- Dust and paint chips from old paint
- Homes built before 1950, particularly those that are in need of repair or are in deteriorating condition
- Soil that has lead in it
- Hobby materials such as stained glass, paints, solders, fishing weights, and buckshot
- Folk remedies
- Workplace dust brought home on the clothing of people who have jobs that use lead, such as battery manufacturers or smelting companies

- Food stored in some ceramic dishes (especially if made in another country)
- Older painted toys and antique furniture such as cribs
- Tap water in homes that have lead pipes
- Mini-blinds manufactured outside the United States before July 1996

Prevention—what you can do

- If your home was built before 1950, ask your pediatrician to test your child for lead.
- If your home was built before 1978, talk to your pediatrician or health department about safe ways to remodel *before* any work is done.
- Know your state's laws regarding lead removal. Some states do not allow home owners to remove lead, only certified de-leaders.
- Clean and cover any chalking, flaking, or chipping paint with a new coat of paint, duct tape, or contact paper. It is important to check for paint dust or flaking paint at window areas where children often play.
- Repair areas where paint is dusting, chipping, or peeling before placing cribs, playpens, beds, or highchairs next to them.
- Encourage your children to wash their hands frequently, especially before eating.
- Check your home or apartment for possible lead contamination before moving in. Keep in mind that landlords are legally responsible for removing any lead found on their property.
- If you work around lead or have hobbies that involve lead, change clothes and shoes before entering your home. Keep clothes at work or wash work clothes as soon as possible.
- Check with your pediatrician or health department to see if your area has a problem with lead in the water.
- If you have lead pipes, run the first morning tap water for 2 minutes before using it for drinking or cooking. Do not use hot tap water for mixing formula, drinking, or cooking.

You can also reduce the risks of lead by making sure your child eats a well-balanced diet. Give your child nutritious, low-fat foods that are high in calcium and iron, like meat, beans, spinach, and low-fat dairy products. Calcium and iron in particular reduce the amount of lead absorbed by the body.

Lead screening

The only way to know for sure if your child has been exposed to lead is to have your pediatrician test your child's blood. Lead screening tests use either a small amount of blood from a finger prick or a larger sample of blood from a vein in the arm. These tests measure the amount of lead in the blood.

Treatment

For children with *low* levels of lead in their blood, identify and eliminate the sources of lead to avoid future health problems. Children with *high* levels of lead in their blood usually need to take a drug that binds the lead in the blood and helps the body get rid of it. This treatment is often done in the hospital and usually is given as a series of shots. Some children with lead poisoning need more than one type of treatment and several months of close follow-up. If the damage is severe, the child may need special schooling and therapy.

Most young children put things other than food into their mouths. They chew on toys, taste the sand at the park, and eat cat food if given the chance. This rarely causes any harm, as long as poisons and sharp objects are kept out of reach. Lead, however, can be very dangerous to children. Infants and toddlers can get sick by putting their fingers in their mouths after touching lead dust, eating lead paint chips, or breathing in lead dust. Lead poisoning can cause learning disabilities, behavioral problems, anemia, or damage to the brain and kidneys. Talk to your pediatrician about getting a blood test, especially if your child is under 3 years of age. Take the steps listed in this brochure to make sure your child does not come into contact with lead.

The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

From your doctor

Should my child be screened for lead?

If you can answer "yes" to any of the following questions, especially numbers 1, 2, and 3, your child may need to be screened for lead. Talk to your pediatrician about lead screening for your child.

1. Does your child live in or regularly visit a house that was built before 1950? This includes a home child care center or the home of a relative.
2. Does your child live in or regularly visit a house built before 1978 that has been remodeled in the last 6 months? Are there any plans to remodel?
3. Does your child have a brother, sister, housemate, or playmate who is being treated for lead poisoning?
4. Have you ever been told that your child has high levels of lead in his or her blood or lead poisoning?
5. Does your child live with an adult whose job or hobby involves exposure to lead?
6. Does your child live near an active lead smelter, battery recycling plant, or other industry likely to release lead into the environment?
7. Does your child live within one block of a major highway or busy street?
8. Do you use hot tap water for cooking or drinking?
9. Has your child ever been given home remedies (azarcon, greta, pay looah)?
10. Has your child ever lived outside the United States?
11. Does your family use pottery or ceramics for cooking, eating, or drinking?
12. Have you seen your child eat paint chips?
13. Have you seen your child eat soil or dirt?
14. Have you been told your child has low iron?

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

The American Academy of Pediatrics is an organization of 57,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of infants, children, adolescents, and young adults.

American Academy of Pediatrics
PO Box 747
Elk Grove Village, IL 60009-0747
Web site — <http://www.aap.org>

Copyright ©1998 American Academy of Pediatrics