



Acetaminophen (Tylenol)

NOTE: An infant's weight needs to be exact for the dosing to be accurate. Determine your child's weight before giving any medication. Contact your pediatrician if you have any questions.

Do not exceed 5 doses in 24 hours	May Give Every	10-11 lbs. 2-3 mo.	12-17 lbs. 4-11 mo.	18-23 lbs. 12-23 mo.	24-35 lbs. 2-3 yr.	36-47 lbs. 4-5 yr.	48-59 lbs. 6-8 yr.	60-71 lbs. 9-10 yr.
-----------------------------------	----------------	-----------------------	------------------------	-------------------------	-----------------------	-----------------------	-----------------------	------------------------

Children's Liquid 160 mg/5 ml	4 - 6 hours	1.25 ml	2.5 ml	3.75 ml	5 ml	7.5 ml	10 ml	12.5 ml
Children's Chewable Tablet 160 mg/tablet	4 - 6 hours	-	-	-	1 tab.	1.5 tab.	2 tab.	2.5 tab.
Suppository 80 mg	4 - 6 hours	-	1 supp.	1 1/2 supp.	2 supp.	-	-	-
Suppository 120 mg	4 - 6 hours	-	-	1 supp.	1 1/2 supp.	-	-	-
Suppository 325 mg	4 - 6 hours	-	-	-	1/2 supp.	3/4 supp.	1 supp.	-

CAUTION: We recommend that you use the dropper or dosage cup that comes with your medicine to avoid overdose. Dosage recommendations on bottles may vary from the recommendation on these pages. We would like you to use our recommended dosages unless your health care provider has given you a different (usually higher!) dosage. Aspirin should not be given to your child.

NOTE: The American Academy of Pediatrics (AAP) recommends that dosages be administered in milliliters (ml).

Friday Night Fever?

The things that make parents worry rarely happen between 9 and 5, but we've got your back 24/7. If you have a healthcare concern, call us first. The call is free and could save you both a sleepless night.

[Click for details](#)



801-922-5200