



Ibuprofen (Advil, Motrin, Nuprin, Medipren)

Do not use more than 4 times a day.

Give every 6-8 hours for fever over 102° rectal	May Give Every	12-17 lbs. 6-11 mo.	18-23 lbs. 12-23 mo.	24-35 lbs. 2-3 yr.	36-47 lbs. 4-5 yr.	48-59 lbs. 6-8 yr.	60-71 lbs. 9-10 yr.
Infant Concentrated Drops 50 mg/1.25 ml*	6-8 hours	1.25 ml	1.875 ml	-	-	-	-
Children's Suspension 100 mg/5 ml	6-8 hours	2.5 ml	3.75 ml	5 ml	7.5 ml	10 ml	12.5 ml
Children's Chewables 100 mg	6-8 hours	-	-	1 tab.	1 1/2 tab.	2 tab.	2 1/2 tab.
Children's Tablets 100 mg	6-8 hours	-	-	-	-	2 tab.	2 tab.

CAUTION: DO NOT GIVE TO AN INFANT UNDER 6 MONTHS OF AGE.

Consult with a physician or pharmacist before combining any medication. Some drugs may contain additional pain relievers. Aspirin should not be given to your child.

NOTE: The American Academy of Pediatrics (AAP) recommends that dosages be administered in milliliters (ml).